



Available online through

www.jbsoweb.com

ISSN 2321 - 6328

Research Article

EVALUATION OF ANTIBACTERIAL ACTIVITY OF ESSENTIAL OILS OF DIFFERENT TYPES *ROSMARINUS OFFICINALIS* L.

Muftah A. Nasib¹, Idress Hamad Attitalla^{1*}, Maraia F. Elmhawi²

¹Omar Al-Mukhtar University, Faculty of Science, Department of Microbiology, Box 919, Al-Bayda, Libya

²Department of Chemistry (Biochemistry), Faculty of Science, Benghazi, University, Benghazi, Libya

*Corresponding Author Email: idressattitalla2004@yahoo.com

Article Received on: 02/10/15 Accepted on: 07/12/15

DOI: 10.7897/2321-6328.03651

ABSTRACT

In this work, we studied antibacterial activity of oils extract of leaves *Rosmarinus officinalis*. Rosemary is well known as a spice and widely used plant in ethno medicine worldwide. This study showed that essential oil extract of *Rosmarinus officinalis* from leaves (grow in Al-Jabal Al Akhdar and house) at 10%, 20% and 30% concentration were effective against both Gram-positive and Gram-negative bacteria, while the essential oil commercial was no effective against both Gram-positive and Gram-negative bacteria at all concentration. We suggest the oils of *Rosmarinus officinalis* (grow in Al-Jabal Al Akhdar and house) rich in phenolic constituent such as α -pinene and 1,8-cineole has the highest anti-oxidant activity against bacterial.

Key words: *Rosmarinus officinalis* L, Essential oil, Bacterial and Antibacterial Activity

INTRODUCTION

Herbal medicines are widely used all over the world. They are often perceived as being natural and therefore harmless. Many herbal remedies individually or in combination with different formulations such as leaf, powder, pastes, decoction, infusion, etc. had been recommended to treat various diseases. Many, if not most of medicinal plants contain flavonoids; such compounds have been associated with several beneficial effects such as anti-oxidation which consider to be a fundamental property important for life¹.

Many plant products exert antioxidant effect by quenching various free radicals and the singlet form of molecular oxygen. Spices exhibit wide range of beneficial, pharmacological, antioxidant, anti-carcinogenic and anti-inflammatory, antimicrobial effects².

Rosemary is a powerful herb belonging to the family Lamiaceae that originates from the Mediterranean region. It is derived from the Latin word *ros* (dew) and *marinus* (sea) which means 'dew of the sea'³. Rosemary (*Rosmarinus officinalis* L.) is a spice and medicinal herb widely used around the world. Of the natural antioxidants, rosemary has been widely accepted as one of the spices with the highest antioxidant activity⁴. Rosemary leaves have many traditional uses based on their antibacterial and spasmolytic actions. Used orally for the treatment of dyspeptic complaints⁵.

Essential oils are natural, concentrated, volatile aromatic compounds isolated from plants. These compounds possess a wide spectrum of pharmacological activities. The main advantage of natural agents is that they do not enhance the "antibiotic resistance", a phenomenon encountered with the long term use of

synthetic antibiotics. One of the known plants that used in the folk medicine is *Rosmarinus officinalis*⁶. Many compounds have been isolated from rosemary, including flavones, diterpenes, steroids, and triterpenes. Of these, the antioxidant activity of rosemary extracts has been primarily related to two phenolic diterpenes: carnosic acid and carnosol⁶. Rosemary essential oil is also used as an antibacterial, antifungal⁷⁻⁸ and anticancer agent⁹.¹⁰ investigated the antibacterial activity of selected essential oils against some food spoilage organisms. They concluded that the essential oils of cinnamon, clove and rosemary were the most active. Similar results were obtained by¹¹ for the antibacterial activity of rosemary essential oil against *Bacillus cereus* strains grown in carrot broth. The purpose of the present work is to determine the antimicrobial activity of essential oils of *Rosmarinus officinalis* grow in Al-Jabal Al Akhdar, and house in Benghazi and commercial from a super market.

MATERIALS AND METHODS

Plant material: The leaves of *Rosmarinus officinalis*, were collected from Al-Jabal Al Akhdar area in Benghazi, Libya 2013 (sample 1), and The leaves of *Rosmarinus officinalis* were collected from house in Benghazi, Libya 2013 (sample 2), but oil of *Rosmarinus officinalis* commercial from a super market, Benghazi, Libya 2013(sample3).

Bacteria used: Bacteria were taken from the laboratory of microbiology in Benghazi medical center, which know as multi drug resistant bacteria. The bacteria used were *Escherichia coli* (MDR)ATCC, *Staphylococcus aureus* (MDR)ATCC, *Pseudomonas aeruginosa* (MDR) ATCC, *Klebsiella pneumonia* (MDR) and *Acinetobacter sp.* (MDR). Other bacteria were not multi drug resistant such as *Escherichia coli*, *Staphylococcus aureus* and *Pseudomonas aeruginosa*. The organisms were

isolated and identified by standard methods, and identification confirmed by using phonex. The organisms were then sub cultured and maintained on nutrient agar slants.

Sample preparation

Extraction of essential oil: The dry powdered leaves of *Rosmarinus officinalis* (sample 1 and sample 2) (500g) were subjected to hydro distillation using Clevenger apparatus. The isolation of volatile oils was complete within 6 hours¹². The oil samples were stored at 7°C in dark air-tight.

Antibacterial activities of essential oil extract rosemary collected from Al-Jabal Al Akhdar, house and commercial rosemary oil: In this study diluted essential oils were used. The diluted essential oils were prepared by using Dimethyl Sulfoxide (DMSO) to obtain 10%(v/v), 20%(v/v) and 30%(v/v) concentrations. DMSO was used as negative control. A screening assay using well diffusion¹³. Muller Hinton agar plates were inoculated by rubbing sterile cotton swabs after immerse 100µ l bacterial suspensions on plates (overnight cultures grown at 37°C on nutrient agar and adjusted to 0.5 McFarland in sterile saline) over the entire surface of the plate. After inoculation 9 mm diameter wells were cut into the surface of the agar using a sterile cork borer. Different concentrations (10, 20 and 30 %) were added to the wells. Plates were incubated at 37°C for 24 h. Control wells contained solvent DMSO. Zones of inhibition were measured by using ruler. The diameter of zones was recorded. Each assay was carried out in triplicate. The antibacterial assay plates were incubated at 37°C for 24hr. The effect of essential oils on the tested bacteria was compared with the sensitivity of the same bacteria to five antibiotics Colisti sulphate, Amicacin, Amoxycillin, gentamicin and sulphamethoxazole trimethoprim (60µg/ml)¹⁴⁻¹⁵

RESULTS

Antibacterial evaluation of essential oil extract rosemary collected from Al-Jabal Al Akhdar, house and commercial rosemary oil

The antibacterial activity of essential oil extract from rosemary collected from Al-Jabal Al Akhdar, house and commercial rosemary oil were tested using different bacteria and the results are listed in Table 1. The effect of essential oils on the tested bacteria was compared with the sensitivity of the same bacteria to five antibiotics Colisti sulphate, Amicacin, Amoxycillin, gentamicin and sulphamethoxazole trimethoprim are shown in Table 2.

This results showed that two type oils extracts of rosemary (collected from Al-Jabal Al Akhdar and house) were effective against both Gram-positive and Gram-negative bacteria, while commercial oil rosemary no any activity against both Gram-positive and Gram-negative bacteria.

It was found that the essential oil from rosemary collected from Al-Jabal Al Akhdar was potentially active against *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Acinetobacter sp.* and *Klebsiella pneumonia* with zones of inhibition ranging from 4 to 9 mm at 10 %, 7 to 12 mm at 20 % and 9 to 18mm at 30%. *Staphylococcus aureus* bacteria showed a better inhibitory effect on the essential oil from rosemary collected from Al-Jabal Al Akhdar at 10 % concentration (9mm) out of five bacterial strains tested, while at 20 % and 30 % concentrations (11mm and 14mm) as shown in figures 5. Figure

2 show the essential oil from rosemary collected from Al-Jabal Al Akhdar inhibit the growth of *E.coli* bacteria strains produced a zone diameter of inhibition from 4 - 11 mm at 10 %,20 %, and 30 % concentrations. Antibacterial activity of essential oil from rosemary collected from Al-Jabal Al Akhdar against *Klebsella.pneumonia* bacteria at 10 %, 20 % and 30 % concentrations (7mm, 12mm and 18mm) respectively are shown in figure 3. The essential oil from rosemary collected from Al-Jabal Al Akhdar showed an inhibitory effect on *P.aeruginosa* bacterial strains with the inhibition zone ranging from 5 to 9 mm at 10 %, 20 % and 30 % concentrations as can be seen in figure 4. The results showed that the essential oil from rosemary collected from Al-Jabal Al Akhdar has on inhibition of *Acinetobacter sp.* bacteria when the concentration 10 % (8mm), while 20 % and 30 % concentrations have the effect (11 and 14mm respectively) as shown in figure 1.

The results showed an inhibition slightly active against *Acinetobacter sp.* at 20% and 30% concentrations of essential oil extracted from rosemary collected from house while no inhibition was observed against *Acinetobacter sp.* at 10% contraction as shown in figures 6.

Figure 7 show the essential oil from rosemary collected from house inhibit the growth of *Klebsella.pneumonia* bacteria strains produced a zone diameter of inhibition from 5 - 9 mm at 10%,20%, and 30% concentrations. Antibacterial activity of essential oil from rosemary collected from house against *Pseudomonas aeruginosa* bacteria at 10%, 20% and 30% concentration (1mm, 8mm and 9mm) respectively are shown in figure 8.

The commercial rosemary oil was not suitable for testing the antimicrobial activity against *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Acinetobacter sp.* and *Klebsiella pneumonia* at all contractions as shown in table 1 and figures 9,10,11,12 and 13.

DISCUSSION

There is an increasing interest in phytochemicals as new sources of natural antioxidant and antimicrobial agents. The use of synthetic antioxidants in the food industry is severely restricted as to both application and level¹⁶. Currently, there is a strong debate about the safety aspects of chemical preservatives, since they are considered responsible for many carcinogenic and teratogenicity attributes, as well as residual toxicity¹⁷

Plant-derived polyphenols receive considerable attention because of their potential antioxidant and antimicrobial properties¹⁷. Phenolic compounds exhibit a considerable free-radical scavenging (antioxidant) activity, which is determined by their reactivity as hydrogen- or electron- donating agents, the stability of the resulting antioxidant-derived radical, their reactivity with other antioxidants and, finally, their metal chelating properties¹⁸.

Del Campo *et al* reported that rosemary plants are rich sources of phenolic compounds with high antimicrobial activity against both Gram-positive and Gram-negative bacteria. High percent of the antimicrobial activity they attributed to carnosic acid and carnosol. It is clear that rosemary extracts have bioactive properties, but their antimicrobial activities have not been deeply characterized¹⁹.

Many compounds have been isolated from rosemary, including flavones, diterpenes, steroids, and triterpenes. Of these, the

antioxidant activity of rosemary extracts has been primarily related to two phenolic diterpenes: carnosic acid and carnosol⁸.

The activity of rosemary has been ascribed to the diterpene content, mainly carnosic acid and carnosol²³, as well as to the essential oil constituents²⁴. Carnosic acid provides protection from the liver carcinogen aflatoxin²⁵.

Rosemary extracts are widely used in the food. Their major bioactive components have shown antioxidant, antimicrobial, anti-inflammatory, anti-tumorigenic and chemo preventive activities²⁵.

The main compounds responsible for the antimicrobial activity are α -pinene, bornyl acetate, camphor and 1,8-cineole²⁰⁻²²

CONCLUSION

This results showed that two type oils extracts of rosemary (collected from Al-Jabal Al Akhdar and house) were effective against both Gram-positive and Gram-negative bacteria, while commercial oil rosemary no any activity against both Gram-positive and Gram-negative bacteria. We suggest the two type oils extracts of rosemary (collected from Al-Jabal Al Akhdar and house) more rich in phenolic constituent such as α -pinene, β -pinene, 1,8-cineol, camphene, limonene and camphor than commercial oil. The phenolic constituent has the highest antioxidant activity against bacterial.

Table 1: Screening of Antibacterial activity of essential oil extracted of rosemary collected from Al-Jabal Al Akhdar, house and commercial rosemary oil

S/No.	Treatment	Concentration (%)	Zone of Inhibition (mm) \pm Standard deviation				
			EC	PA	SA	KP	AS
1.	C	-	-	-	-	-	-
2.	essential oil extracted of rosemary collected from Al-Jabal Al Akhdar	10	4 \pm 0.31	5 \pm 0.01	9 \pm 0.15	7 \pm 0.07	8 \pm 0.11
		20	7 \pm 0.26	7 \pm 0.24	11 \pm 0.24	12 \pm 0.09	11 \pm 0.21
		30	11 \pm 0.22	9 \pm 0.12	14 \pm 0.07	18 \pm 0.27	14 \pm 0.09
3.	essential oil extracted from rosemary collected from house	10	9 \pm 0.31	1 \pm	9 \pm 0.01	5 \pm 0.03	-
		20	14 \pm 0.26	8 \pm 0.33	11 \pm 0.04	7 \pm 0.15	2 \pm 0.35
		30	20 \pm 0.22	9 \pm 0.92	14 \pm 0.21	9 \pm 0.21	2 \pm 0.21
	commercial rosemary oil	10	-	-	-	-	-
		20	-	-	-	-	-
		30	-	-	-	-	-

Values are expressed as Mean (X) \pm SD, n=3 **Abbr.** EC= *Escherichia coli*, PA = *Pseudomonas aeruginosa*, KP = *Klebsiella pneumonia*, SA= *Staphylococcus aureus*, AS= *Acinetobacter sp.*, C: Control (DMSO).

Table 2: Antibiotic activity of different type of bacter

Antibiotic	Zone of Inhibition (mm) \pm Standard deviation				
	EC	SA	PA	AS	KP
Colisti sulphate	-	2 \pm 0.01	3 \pm 0.01	6 \pm 0.03	4 \pm 0.01
Amicacin	15 \pm 0.02	13 \pm 0.02	9 \pm 0.01	-	12 \pm 0.04
Amoxycillin	-	3 \pm 0.01	-	-	2 \pm 0.01
Gentamycin	10 \pm 0.03	6 \pm 0.01	5 \pm 0.02	3 \pm 0.03	1 \pm 0.01
Sulphmethoxazole	3 \pm 0.12	19 \pm 0.03	-	4 \pm 0.08	-

Values are expressed as Mean (X) \pm SD, n=3. EC= *Escherichia coli*, PA = *Pseudomonas aeruginosa*, KP = *Klebsiella pneumonia*, SA= *Staphylococcus aureus*, AS= *Acinetobacter sp.*



Figure (1): Antibacterial activity of essential oils extracted of leaves grown in Al-Jabal Al Akhdar against *Acinetobacter sp* bacteria at 10 % ,20 % and 30 % concentrations




Figure (2): Antibacterial activity of essential oils extracted of leaves grown in Al-Jabal Al Akhdar against *E.coli* bacteria at 10 % ,20 % and 30 % concentrations




Figure (3): Antibacterial activity of essential oils extracted of leaves grow in Al-Jabal Al Akhdar against *Klebsella.pneumonia* bacteria at 10 % ,20 % and 30 % concentrations




Figure (4): Antibacterial activity of essential oils extracted of leaves grow in Al-Jabal Al Akhdar against *P.aeruginosa* bacteria at 10 % ,20 % and 30 % concentrations



Figure (5): Antibacterial activity of essential oils extracted of leaves grow in Al-Jabal Al Akhdar against *Staphylococcus.arues* bacteria at 10 % ,20 % and 30 % concentrations




Figure (6): Antibacterial activity of essential oils extracted of leaves grow in house in Benghazi from *Rosmarinus officinalis* against *Acinetobacter sp.* bacteria at 10% ,20% and 30% concentrations



Figure (7): Antibacterial activity of essential oils extracted of leaves grow in house in Benghazi from *Rosmarinus officinalis* against *Klebsella.pneumonia* bacteria at 10% ,20% and 30% concentrations




Figure (8): Antibacterial activity of essential oils extracted of leaves grow in house in Benghazi from *Rosmarinus officinalis* against *P.aeruginosa* bacteria at 10% ,20% and 30% concentrations




Figure (9): Antibacterial activity of commercial rosemary oil against *Acinetobacter sp.* bacteria at 10% ,20% and 30% concentration




Figure (10): Antibacterial activity of commercial rosemary oil against *E.coli* bacteria at 10% ,20% and 30% concentration




Figure (11): Antibacterial activity of commercial rosemary oil against *Klebsella pneumonia* bacteria at 10% ,20% and 30% concentration




Figure (12): Antibacterial activity of commercial rosemary oil against *Pseudomonas aeruginosa* bacteria at 10% ,20% and 30% concentration



Figure (13): Antibacterial activity of commercial rosemary oil against *Staphylococcus aureus* bacteria at 10% ,20% and 30% concentration

ACKNOWLEDGEMENT

Authors are thankful Dr. Fatima A. Elslimani in Microbiology Department in Benghazi Medical Center - Libya for carrying the present work.

REFERENCES

- Halliwell, B. and Gutteridge, J.M.C. The importance of free radicals and catalytic metal ions in human diseases. *Mol. Asp. Med.*2010; 8: 89-93.
- Sachin Uttam Rakesh, Priyanka R. Patil, and Sagar R. Mane (2010): Use of Natural Antioxidants to Scavenge Free Radicals: A Major Cause of Diseases, *International Journal of Pharm Tech Research.*2010; 2(2): 1074-1081.
- All about rosemary (Rosmarinus officinalis). (2012). <http://www.apinchof.com/rosemary1066.html>
- Peng, Y., Yuan, J., Liu Fand, J. Y. Determination of active components in rosemary by capillary electrophoresis with electrochemical detection. *Journal of Pharmaceutical and Biomedical Analysis.*2005; 39, 431-437.
- Ho C-T, Ferraro T, Chen Q, Rosen RT, Huang MT. Phytochemicals in teas and rosemary and their cancer preventive properties. American chemical society. Washington DC, 1994;2-19 Zorica Stojanović-Radić, Marija Nešić, Ljiljana Čomić, Niko Radulović Antimicrobial activity and cytotoxicity of commercial rosemary essential oil (Rosmarinus officinalis L.). *biologica nyssana.*2010; (1): 83-88.
- Oluwatuyi M, Kaatz GW, Gibbons S. Antibacterial and resistance modifying activity of Rosmarinus officinalis. *Phytochemistry.*2004; 65: 3249–3254.
- Frankel, E., N. Patil ,R. and Sagar R. Mane . Antioxidant activity of a rosemary extract and its constituents, carnosic acid, carnosol, and rosmarinic acid, in bulk oil and oil-in-water emulsion. *Journal of Agricultural and Food Chemistry.*1996;44(1):. 131-135.
- Leal, P. F. et al. (2003): Functional Properties of Spice Extracts Obtained via Supercritical Fluid Extraction. *Journal of Agricultural and Food Chemistry.*2003; 51(9): 2520-2525.
- Sawsan S. Sayd, Hanan A.A. Taie, Lobna S. Taha. Micropropagation, anti-oxidant activity, total phenolics and flavonoids content of gardenia jasminoides ellis as affected by growth regulators. *International journal of academic research.* 2010; 2: (3). 184-191.
- Himaja Malipeddi, Kalpana Jayapalu, Chaitra Muralidhara. Validated zero and first order derivative spectrophotometric methods for the estimation of poorly soluble antiretrovirals. *Int. J. Res. Ayurveda Pharm.* 2015;6(3):366-370 <http://dx.doi.org/10.7897/2277-4343.06371>.
- Maraia F. Elmhdwi and Idress Hamad Attitalla. Antioxidant and free radical scavenging activity of essential oil extracted from Rosmarinus Officinalis L. *J Bio Sci Opin.* 2015; 3(5):223-229 DOI 10.7897/2321-6328.03549
- Tavassoli ,S. and Emam Djomeh ,Z. Total Phenols, Antioxidant Potential and Antimicrobial Activity of Methanol Extract of Rosemary (Rosmarinus officinalis L.) *Global Veterinaria.* 2011; 7 (4): 337-341.
- Ruberto, G., and Barata, M.T. Antioxidant activity of selected essential oil components in two lipid model systems. *Food Chem.* 2012; 69: 167–174.
- Oyedeki OA, Adeniyi BA, Ajayi O, Konig WA . Essential oil composition of *Piper guineense* and its antimicrobial activity. Another Chemotype from Nigeria. *Phytother Res.* 2005;19:362-364.
- Aziza K.G., Haiko H., Smânia A.Jr, Machado de Souza S. Rosemary (Rosmarinus officinalis) – a study of the composition, antioxidant and antimicrobial activities of extracts obtained with supercritical carbon dioxide. *Cienc. Technol. Aliment., Campinas.* 2008;28, 2, 463-469.
- Mallya Suma V. Potent antioxidant from a traditional herbal heritage. *J. Biol. Sci. Opin.* 2015;3(2):57-60 <http://dx.doi.org/10.7897/2321-6328.03213>
- Nyangono, C.F., Chakokam Ngangoum, R.M, Kuate,I.D, Ngondi,J.L, Enyong Oben, J. Effect of Guibourtia tessmannii extracts on blood lipids and oxidative stress markers in triton WR 1339 and high fat diet induced hyperlipidemia rats. *Biology and Medicine.* 2012; 4 (1): 01- 09.
- Del Campo, J., M.J. Amiot and C. Nguyen (2000): The Antimicrobial effect of rosemary extracts. *J. Food Protect.* 2000; 10: 1359-1368.
- Daferera, D. J.; Ziogas, B. N.; Polissiou, M. G . The effectiveness of plant essential oils in the growth of Botrytis cinerea, Fusarium sp. and Clavibacter michiganensis subsp. michiganensis. *Crop Protection.* 2003; 22(1):39-44.
- Daferera, D. J.; Ziogas, B. N.; Polissiou, M. G GC-MS analysis of essential oils from some Greek aromatic plants and their fungitoxicity on Penicillium digitatum. *Journal of Agricultural and Food Chemistry.*2000; 48(6): 2576-2581.
- Pintore, G., Usai, M., Bradesi, P., Juliano, C., Boatto, G., Tomi, F., Chessa, M., Cerri, R., Casanova, J. Chemical composition and antimicrobial activity of Rosmarinus officinalis L. oil from Sardinia and Corsica. *Flav. Fragr. J.* 2002;17: 15-19
- Asia Begum, Subarda Sandhya, Syed Shaff ath Ali, Kombath Ravindran Vinod, Swapna Reddy, David Banji An in-depth review on the medicinal flora rosmarinus officinalis (lamiaceae). *Acta Sci. Pol., Technol. Aliment.*2013; 12(1): 61-73
- Asia B., Subarda S., Syed Shaff A., Kombath R., Vinod S., David B. An in-depth review on the medicinal flora rosmarinus officinalis (lamiaceae). *Acta Sci. Pol., Technol. Aliment.*2013; 12(1), 61-73
- Gutiérrez R., Alvarado J.L., Presno M., Perez-Veyna O., Serrano C.J., Yahuaca P. Oxidative stress modulation by Rosmarinus offi

- cinalis in CCl₄-induced liver cirrhosis. *Phytother. Res.* 2010; 24 (4):595-601.
25. Bozin B, Mimica-Dukic N, Samojlik I and Jovin E (2007): Antimicrobial and antioxidant properties of rosemary and sage (*Rosmarinus officinalis* L. and *Salvia officinalis* L., Lamiaceae) essential oils. *J Agric Food Chem.* 2007; 55(19): 7879–7885.

Cite this article as:

Muftah A. Nasib, Idress Hamad Attitalla, Maraia F. Elmhawi. Evaluation of antibacterial activity of essential oils of different types *Rosmarinus officinalis* L. *J Biol Sci Opin* 2015;3(6):235-240
<http://dx.doi.org/10.7897/2321-6328.03651>

Source of support: Nil; Conflict of interest: None Declared

Disclaimer: JBSO is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the contents published in our Journal. JBSO cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of JBSO editor or editorial board members.